arXiv:1702.03448v1 [hep-th] 11 Feb 2017

February 14, 2017

Ultra-Spinning Limit of Charged Kaluza-Klein AdS Black Holes and Their 2d CFT Duals

S.M. Noorbakhsh[1] **and M. Ghominejad**[2]

Department of Physics, Semnan University, P.O. Box 35195-363, Semnan, Iran

Abstract

By using ultra-spinning limit as a generating solution technique, we construct a new class of charged rotating Kaluza-Klien AdS black holes. That describes the exact solutions of Einstein-Maxwell dilaton theory in all higher dimensions. The obtained geometries possess some punctures, describing a noncompact horizon, but has a finite area. We then explicitly investigate the validity of the Kerr/CFT correspondence for the four-dimensional case. Also, for the higher-dimensional solutions we find a main result for the central charges. Then via the Cardy's formula we argue the existence of a precise agreement between the microscopic entropy of the dual CFT and the entropy of the noncompact horizon.

[1]e-mail: m.noorbakhsh@semnan.ac.ir
[2]e-mail: mghominejad@semnan.ac.ir

1 Introduction

It was shown that within the higher-dimensional general relativity one can access to the rotating black holes with any arbitrary large angular momentum by keeping the mass fixed [1]. In order to study the stability of Myers-Perry black hole in a large angular momentum per unit mass, such that the rotation parameter $a \to \infty$, Emparan and Myers introduced the "ultra-spinning" black holes [2]. The horizon geometry of their resulting solutions have a pancake shape, spreading out along the rotation plane while are contracted in the transverse directions. They also found a limit in which these rapid black holes become an unstable black membrane at a finite large rotation. It was generalized an analogous analysis for asymptotic AdS rotating black holes. In $d \geq 6$ what is called as a black brane ultra-spinning limit proposed in [3] by which the rotation parameter a_i approaches to the AdS radius l, while one has to keep the physical mass and angular momentum finite, thereby leads to a static black brane. Furthermore, a different technique was introduced in [4] that ultra-spinning (A)dS solutions can be derived such that both a and l simultaneously approach to infinity while imposing a/l fixed. Meanwhile, Caldarelli *et al.* studied other applicable technique [3, 5], in which by taking $a \to l$ limit the horizon radius r_+ remains fixed, while the polar angle θ is scaled in a special way to avoid a metric singularity. This limit gives rise to a noncompactness black hole solution with a horizon topology as $\mathbb{H}^2 \times S^{d-4}$. So this ultra-spinning method refers to a hyperboloid membrane limit.

In [6] it was introduced a novel ultra-spinning approach that can be used only in the presence of the negative cosmological constant. The constructing geometries have a noncompact horizon that topologically can be described as a sphere with some punctures, but has a finite area. These class of black holes, in a sense, correspond to those were constructed in $\mathcal{N} = 2$ gauged supergravity in the presence of a scalar potential, which have similar horizon topology [7, 8]

In the ultra-spinning technique introduced [6] to gain a regular metric, one should first transform the given rotating AdS black hole to an asymptotically rotating frame, then before boosting this rotation to the speed of light by taking $a \to l$, one should replace the corresponding azimuthal coordinate by a new scaled proper coordinate. This method has been investigated for the general multi-spinning Kerr-AdS black holes in [9], and for two particular classes of gauged supergravity solutions in [10]. The obtained geometries as a new classes of ultra-spinning solutions exhibit a distinguishable property in the extended thermodynamic phase space point of view, where the variation of the cosmological constant has to be taken in to account [11]. It was shown that the entropy of these specific solutions in some range of parameter space violate the Reverse Isoperimetric Inequality. So, this ultra-spinning limit is denoted as super-entropic limit as well. [6, 9].[1]

There are strong evidences indicating black holes can be viewed as thermodynamic systems possessing underlying microstate structure, attributed to a theory of quantum gravity. To provide an appropriate quantum gravity theory, proposing a successful formalism to calculate the entropy of black holes is one of the most crucial challenge. It was a long historical idea supporting that only microstates near the black hole horizon should be considered to calculate the black hole entropy [13–16]. For extremal black holes, the Kerr/CFT correspondence gives a remarkable observation that the statistical microscopic entropy of the quantum gravity theory exactly matches with the Bekenstein-Hawking entropy of the black holes [17]. The basic idea follows from the fundamental example of

[1] It was shown that for a large class of black holes, for a given thermodynamic volume the maximum entropy is related to the (charged) AdS-Schwarzschild black hole. The 'superentropic' black holes referring to those their entropy exceed the maximal entropy [12].

Brown and Henneaux [18], shown by considering consistent boundary conditions, the asymptotic symmetry group (ASG) of AdS_3 can be generated by two copies of Virasoro algebra. It asserts that the quantum theory of gravity on AdS_3 can be described by a two dimensional conformal field theory. Furthermore, Strominger *et al.* via the Cardys formula [19] found a precise agreement between the Bekenstein-Hawking entropy of the Kerr black holes and entropy of the near-horizon quantum states [14, 20]. Therefore what is dubbed the Kerr/CFT correspondence, conjectures an extreme Kerr black hole is holographically dual to a two-dimensional chiral (left-moving part) conformal field theory. These proposal have been developed to the large classes of higher dimensional black hole solutions in asymptotically flat and AdS spacetime [21–23]. In D dimensional Kerr black holes we are dealing with $[(D-1)/2]$ central charges associated to each commutating Virasoro algebra. That using the Cardy's formula one can reproduce the same black hole area law entropy. Moreover, there are numerous progresses to explore the validity of the Kerr/CFT within the various horizon topology. For instance, for the ultra-spinning solutions with a non-compact horizon, the existence of a well-defined Kerr/CFT is also confirmed by [10] and [24].

In this work, we apply the novel ultra-spinning (super-entropic) limit introducing by [6], to the general charged Kaluza-Klein black holes to generate a new class of rotating black hole solutions in that same theory. The first derivation of four dimensional charged Kaluza-Klein AdS (KK-AdS) black hole constructed in [25] by a dimensional reduction of the boosted Five dimensional neutral Kerr-AdS black hole, that presents an exact solution of four dimensional Einstein-Maxwell-Dilaton theory. Its generalization by including NUT charge obtained in [26]. A general solutions in all higher dimensions were also found in [27]. In the context of gauged supergravity, the charged KK black hole solutions in four and five dimensions were introduced in [28,29]. The general metric of multi-spinning charged KK-AdS black holes in all higher dimensions was constructed by Wu [30], that also describes an exact solution of D dimensional Einstein-Maxwell dilaton theory.

The remarkable properties of ultra-spinning solutions, motivated us to further explore this limit onto the multi spinning charged KK-AdS black holes as a simple generating solution method to construct a new general class of solution. For this purpose we find the general metric of ultra-spinning charged KK-AdS black holes in all dimensions. Then we show the obtained geometries enjoy a noncompact horizon but has a finite area. Furthermore, we check the validity of the Kerr/CFT correspondence for our resulting solutions in all dimensions.

The rest of this paper is organized as follows. In section 2 we introduce the general charged KK-AdS black holes we shall analyse, reviewing their charges and thermodynamic. Then we obtain their general ultra-spinning version in all dimensions, followed by their horizon geometries showing a noncompact manifold. We then present the explicit metric for four dimensional case. In the Sec. 3 we study the ultra-spinning Kerr/CFT description, that in four dimensional case we precisely show that the entropy of the CFT side via the Cardy's formula agrees with the black hole entropy. Then in the last subsection, we find the near horizon geometry of the general ultra-spinning KK-AdS solutions in all dimensions. We also provide a main result confirming these non-compactness horizon solutions exhibit a well-defined Kerr/CFT correspondence.

2 General charged rottaing Kaluza-Klein AdS black holes

Here, we consider the general class of charged multi-spinning KK-AdS black holes, constructed by Wu [30]. That describes a general solution of the Einstein-Maxwell dilaton theory

$$\mathcal{L} = \sqrt{-g}\big\{R - \frac{1}{4}(D-1)(D-2)(\partial\Phi)^2 - \frac{1}{4}\epsilon^{-(D-1)\Phi}F^2 + g^2(D-1)[(D-3)\epsilon^{\Phi} + e^{-(D-3)\Phi}]\big\}. \quad (2.1)$$

The general solution of this theory is given by

$$ds^2 = H^{\frac{1}{D-2}}\bigg[d\gamma^2 + \frac{U dr^2}{\Delta} + \frac{2m}{UH}\omega^2 + d\Omega^2\bigg], \quad (2.2)$$

where we have defined

$$\begin{aligned} d\gamma^2 &= -\frac{W\rho^2}{l^2}dt^2 + \sum_{i=1}^{N}\frac{r^2+a_i^2}{\Xi_i}\mu_i^2 d\phi_i^2, \quad (2.3)\\ d\Omega^2 &= \sum_{i=1}^{N+\varepsilon}\frac{r^2+a_i^2}{\Xi_i}d\mu_i^2 - \frac{1}{W\rho^2}\bigg(\sum_{i=1}^{N+\varepsilon}\frac{r^2+a_i^2}{\Xi}\mu_i d\mu_i\bigg)^2,\\ \omega &= c\,W dt - \sum_{i=1}^{N}\frac{a_i\sqrt{\chi_i}}{\Xi_i}\mu_i^2 d\phi\,, \end{aligned}$$

and

$$\begin{aligned} \rho^2 &= r^2 + l^2, \qquad H = 1 + \frac{2ms^2}{U}, \qquad U = r^{\varepsilon}\sum_{i=1}^{N+\varepsilon}\frac{\mu_i^2}{r^2+a_i^2}\prod_{j=1}^{N}(r^2+a_j^2), \quad (2.4)\\ W &= \sum_{i=1}^{N+\varepsilon}\frac{\mu_i^2}{\Xi_i}, \qquad F = \frac{l^2r^2}{\rho^2}\sum_{i=1}^{N+\varepsilon}\frac{\mu_i^2}{r^2+a_i^2}, \qquad f(r) = c^2 - s^2\rho^2/l^2,\\ \Delta &= \frac{r^{\varepsilon-2}\rho^2}{l^2}\prod_{i=1}^{N}(r^2+a_i^2) - 2mf(r), \qquad \chi_i = c^2 - s^2\Xi_i, \qquad \Xi_i = 1 - a_i^2/l^2. \end{aligned}$$

Also we have $c = \cosh\delta$ and $s = \sinh\delta$. Now we should notice that to present the general solution above for any dimension D, one needs to define the following conventions

$$d = 2N + 1 + \varepsilon, \quad (2.5)$$

where one set $\epsilon = 0$ and 1 for even and odd dimensions respectively, by requiring $a_{N+1} = 0$ for even dimensions. Also $N = [(D-1)/2]$ denotes the number of azimuthal directions ϕ_i corresponding to N independent rotation parameters a_i with periodicity 2π. Moreover, there are $[D/2]$ numbers of "direction cosines" μ_i's as the remaining spatial coordinates subject to the following constraint

$$\sum_{i=1}^{N+\varepsilon}\mu_i^2 = 1. \quad (2.6)$$

The gauge field is also given by

$$A = \frac{2ms}{U\,H}\big(c\,W dt - \sum_{i=1}^{N}\frac{a_i\sqrt{\chi_i}}{\Xi_i}\mu_i^2 d\phi\big), \quad (2.7)$$

Note, the metric (2.2) is written in an asymptotically static frame (ASF). In the uncharged case ($\delta = 0$), this solution reduces to the general class of Kerr-AdS black holes presenting in [31] and [32]. Also in the case of vanishing cosmological constant, the metric (2.2) corresponds to those introduced in [33]. Particularly, the supergravity solutions derived in [34] by only one charge are equivalent with a nonrotatting case of the metric above in D= $4, 5, 7$ dimensions.

Charges and Thermodynamics: The KK-AdS metric (2.2) describes a black hole, if one can find a real root from the equation $\Delta = 0$. The outer root (r_+) describes a killing horizon is generated by the killing vector$K = \partial_t + \sum_i^N \Omega_i \partial_{\phi_i}$. where Ω_i's denote the angular velocity on the horizon in the ASF.

For the metric (2.2) the Hawking temperature and entropy are given by

$$T = \frac{\sqrt{f(r_+)}[V'(r_+) - 2mf'(r_+)]}{4\pi^2 r_+^{\varepsilon-2} c \prod_{i=1}^N (r_+^2 + a_i^2)}, \qquad S = \frac{\mathcal{V}_{D-2}\, m\, r_+\, c\, l^2 \sqrt{f(r_+)}}{2\rho_+^2) \prod_{i=1}^N \Xi_i}, \tag{2.8}$$

where $\mathcal{V}_{D-2}$ refers to the volume of the unit $(D-2)$-sphere as

$$\mathcal{V}_{D-2} = \frac{2\pi^{(D-1)/2}}{\Gamma[(D-1)/2]}. \tag{2.9}$$

Also the angular velocity and the electric potential on the horizon are

$$\Omega_i = \frac{\rho_+^2 a_i \sqrt{\chi_i}}{c\, l^2 (r_+^2 + a_i^2)}, \qquad \Phi = \frac{s}{c\, l^2} \rho_+^2. \tag{2.10}$$

The Mass, angular momenta and electric charge of this family of black hole solutions are calculated in [**?**], as follows

$$\begin{aligned} M &= \frac{\mathcal{V}_{D-2}\, m}{8\pi \prod_{j=1}^N \Xi_j} \left[c^2 \left(\sum_{i=1}^N \frac{2}{\chi_i} + \varepsilon - 2 \right) + 1 \right], \\ J_i &= \frac{\mathcal{V}_{D-2}\, m a_i c \sqrt{\chi_i}}{4\pi \Xi_i \prod_{j=1}^N \Xi_i}, \qquad Q = \frac{(D-3)\mathcal{V}_{D-2}\, mcs}{8\pi \prod_{j=1}^N \Xi_j}. \end{aligned} \tag{2.11}$$

It is shown in [30] that the charges above satisfy the first law of thermodynamics

$$dM = TdS + \sum_{i=1}^N \Omega_i dJ_i + \Phi dQ, \tag{2.12}$$

2.1 Ultra-Spinning limit

In what follows we shall generate a new class of ultra-spinning charged KK-AdS black holes based on the novel ultra-spinning technique introduced in [6]. This ultra-spinning method can be utilized by following three steps, i) to achieve a non-singular metric, one should transform the given rotating AdS black hole to an asymptotically rotating frame. Then the corresponding azimuthal coordinate should be scaled by a new coordinate transformation. ii) boosting the selected rotation to the speed of light easily by taking the limit $a \to l$. iii) finally, compactifying the new azimuthal coordinate.

At the beginning, towards producing an ultra-spinning black hole upon the metric (2.2), we choose a certain φ_j coordinate as an ultra-spinning direction. Before applying next steps, we start

by a trick used in [9] for separating the ϕ_j direction and taking the limit $a_j \to l$. Thus, the following important result can be derived

$$W\Xi_j \to \mu_j^2. \tag{2.13}$$

Now using the relation above, $d\Omega^2$ becomes

$$d\Omega^2 \;\to\; d\Omega_s^2 = \sum_{i\neq j}^{N+\varepsilon} \frac{r^2+a_i^2}{\Xi_i} d\mu_i^2 - 2\frac{d\mu_j}{\mu_j}\Big(\sum_{i\neq j}^{N} \frac{r^2+a_i^2}{\Xi_i}\mu_i d\mu_i\Big) + \frac{d\mu_j^2}{\mu_j^2}\big(\rho^2\hat{W} + l^2\mu_j^2\big), \tag{2.14}$$

where $\hat{W} = \sum_{i\neq j}\frac{\mu_i^2}{\Xi_i}$. Now we are ready to perform the ultra-spinning limit. So in the first step we have to use the following coordinate transformation to gain an asymptotically rotating frame (ARF),

$$\phi_j = \phi_j^R + \frac{a_j}{l^2}t. \tag{2.15}$$

so, we get

$$\omega = (c\hat{W} + c\frac{\mu_j^2}{\Xi_j} - \frac{a_j^2\sqrt{\chi_i}}{l^2\Xi_j}\mu_j^2)dt - \frac{a_j\mu_j^2\sqrt{\chi_i}}{\Xi_j}d\phi_j^R - \sum_{i\neq j}\frac{a_i\sqrt{\chi_i}}{\Xi_i}\mu_i^2 d\phi_i. \tag{2.16}$$

Now, by replacing the new coordinate transformation

$$\varphi_j = \frac{\phi_j^R}{\Xi_j}, \tag{2.17}$$

and upon boosting $a \to l$, we find the new geometry as

$$ds^2 \;=\; \hat{H}^{\frac{1}{D-2}}\left[d\gamma_s^2 + \frac{\hat{U}dr^2}{\hat{\Delta}} + \frac{2m}{\hat{U}\hat{H}}\omega_s^2\right] + d\Omega_s^2, \tag{2.18}$$

where

$$\begin{aligned}
w_s &= (c\hat{W} + \frac{2+3s^2}{2c}\mu_j^2)dt - c\,l\mu_j^2 d\varphi_j - \sum_{i\neq j}^{N}\frac{a_i\mu_i^2 d\phi_i}{\Xi_i},\\
d\gamma_s^2 &= -\Big(\rho^2(\hat{W}+\mu_j^2) + \mu_j^2 l^2\Big)\frac{dt^2}{l^2} + \rho^2\mu_j^2 d\varphi_j^2 + \frac{2\rho^2\mu_j^2 dt d\varphi}{l} + \sum_{i\neq j}\frac{r^2+a_i^2}{\Xi_i}\mu_i^2 d\phi_i^2,\\
\hat{U} &= r^{\varepsilon}\Big(\mu_j^2 + \sum_{i\neq j}\frac{\mu_j^2\rho^2}{r^2+a_j^2}\Big)\prod_{k\neq j}^{N}(r^2+a_k^2), \qquad \hat{F} = \frac{r^2l^2}{\rho^2}\Big(\frac{\mu_j^2}{\rho^2} + \sum_{i\neq j}\frac{\mu_i^2}{r^2+a_i^2}\Big),\\
\hat{\Delta} &= \frac{r^{\varepsilon-2}\rho^4}{l^2}\prod_{i\neq j}^{N}(r^2+a_i^2) - 2mf(r).
\end{aligned} \tag{2.19}$$

At the end, since φ is a noncompact coordinate, one can identify it by requiring that $\varphi \sim \varphi + \mu$.

Also, the gauge field (2.7) under the ultra-spinning limit takes the following form

$$A = \frac{2ms}{\hat{U}\,\hat{H}}\Big((c\hat{W} + \frac{2+3s^2}{2c}\mu_j^2)dt - c\,l\mu_j^2 d\varphi_j - \sum_{i\neq j}^{N}\frac{a_i\mu_i^2 d\phi_i}{\Xi_i}\Big), \tag{2.20}$$

We note that ∂_{φ_j}, is a Killing vector of obtained geometry (2.18). Therefore one can show that the new metric is an exact solution of the theory (2.1). Moreover, the gauge field (2.20) satisfies the equation of motion of this theory as well. The metric (2.2) in the case of vanishing charge $\delta = 0$ reproduces the ultra-spinning general Kerr-AdS black holes investigating in [9].

Basic properties: The new solution (2.18) describes a black hole whenever the equation $\hat{\Delta} = 0$ allows a real root. The largest root (r_+) again can be generated by the Killing vector field

$$K = \partial_t + \Omega_j \partial_{\varphi_j} + \Omega_i \partial_{\phi_i}, \tag{2.21}$$

where Ω_j and Ω_i are the angular velocities on the horizon

$$\Omega_{i\neq j} = \frac{(r_+^2 + l^2) a_i \sqrt{\chi_i}}{c\, l^2 (r_+^2 + a_i^2)}, \qquad \Omega_j = \frac{l^2(s^2+2) - r_0^2 s^2}{2l(s^2+1)(l^2+r_0^2)}. \tag{2.22}$$

Also for the new general solution (2.18), the Hawking temperature can be derived through the horizon surface gravity as

$$T = \frac{\sqrt{f(r_+)}[\hat{V}'(r_+) - 2m f'(r_+)]}{4\pi^2 \rho^2 r_+^{\varepsilon-2} c \prod_{i\neq j}^{N}(r_+^2 + a_i^2)}, \tag{2.23}$$

The electric potential on the horizon is calculated using the definition $\Phi = -K^\mu A_\mu|_{r=r_+}$, leading to

$$\Phi = \frac{s\,(r_+^2 + l^2)}{c\, l^2}. \tag{2.24}$$

Let us here focus on the geometry of the horizon

$$ds_h^2 = \hat{H}^{\frac{1}{D-2}} \left[\rho^2 \mu_j^2 d\varphi_j^2 + \sum_{i\neq j} \frac{r_+^2 + a_i^2}{\Xi_i} \mu_i^2 d\phi_i^2 + \frac{2m}{\hat{U}\hat{H}} \left(c\, l \mu_j^2 d\varphi_j + \sum_{i\neq j}^{N} \frac{a_i \mu_i^2 d\phi_i}{\Xi_i} \right) \right] + d\Omega_s^2. \tag{2.25}$$

At the first glance the $d\Omega_s^2$ seems to be singular at $\mu_j = 0$, but by examining the behavior of the horizon metric near $\mu_j = 0$, one can show that these poles are not true curvature singularity. For simplicity we consider the ϕ_i =const. and μ_i =const. slices. Then the metric above in the leading order expansion of small μ_j takes the following explanation

$$ds_h^2 \approx \hat{H}^{\frac{1}{D-2}} \rho^2 \hat{W} \left[\rho^2 \mu_j^2 d\varphi_j^2 + \frac{2\,m}{\hat{W}\rho^2 \hat{U}\hat{H}} c^2 l^2 \mu_j^4 d\varphi_j^2 + \frac{d\mu_j^2}{\mu_j^2} \right]. \tag{2.26}$$

One can check that this metric present a constant negative curvature geometry on a quotient of the hyperbolic space $\mathbb{H}^2$. Namely, the poles $\mu_j = 0$ are no part of the spacetime but can be interpreted as a kind of boundary. Therefore any geometry of constant (t, r) slices, and particularly the horizon metric are noncompact manifolds, topologically are spheres containing some punctures. However one can find its area and entropy as a finite value

$$S = \frac{\mathcal{V}_{D-2}\, r_+^{\epsilon-1}\, c}{4\sqrt{f(r_+)}} \prod_{i\neq j}^{N} \frac{r_+^2 + a_i^2}{\Xi_i}. \tag{2.27}$$

We should emphasis that it remains to be impossible to obtain a multi ultra-spinning solution. Because taking another coordinate as an ultra-spinning direction, leads the components of $d\Omega_s^2$ to diverge, which can not be removed by another new coordinate transformation.

To better understanding of the obtained geometry (2.18), we shall show the explicit ultra-spinning KK-AdS metric through four dimension in the next subsections.

2.2 Four Dimensional Ultra-Spinning KK-AdS Black Holes

In the four dimensional case, we have only one azimuthal coordinate ϕ corresponding to one rotational parameter a, and there is one spatial polar coordinate θ. So, using the common convention

$$\mu_1 = \sin\theta, \qquad\qquad \mu_2 = \cos\theta, \tag{2.28}$$

the explicit metric (2.18) in 4D takes the following description

$$ds^2 = \sqrt{1+\frac{2mrs^2}{\Delta_\theta}}\Big[\Delta_r dt^2 + \frac{l^2\,\Delta_\theta}{\Delta_r}dr^2 + \frac{2\,mr\,c^2}{\Delta_\theta + 2mrs^2}\big(l\sin^2\theta d\varphi + \frac{\Sigma\,\cos^2\theta}{2m\,l^2\,c^2}dt\big)^2 + \frac{\Delta_\theta}{\sin^2\theta}d\theta^2\Big], \tag{2.29}$$

where

$$\begin{aligned}\Delta_r &= (r^2+l^2)^2 - 2mr(l^2-r^2s^2)2\,m\,rl^4c^2, \qquad\qquad \Delta_\theta = r^2 + l^2\cos^2\theta,\\ \Sigma &= \frac{l^2(r^2+l^2+mrs^2)\cos^2\theta}{r} + 2m(r^2s^2-l^2) + l^2(r-ms^2) + r^3.\end{aligned} \tag{2.30}$$

For this black hole, the hawking temperature, entropy and angular velocity on the horizon read

$$T = \frac{\sqrt{l^2-r_+^2s^2}\,(l^4+r_+^4s^2-3c^2l^2r_+^2)}{4\pi\,c\,l^3r_+(l^2+r_+^2)}, \qquad S = \frac{\mu\,l\,c\,(l^2+r_+^2)}{2(l^2-r_+^2s^2)}.$$

In order to visualize the noncompactness horizon of the metric (2.29), one can embed the horizon metric in an Euclidean 3-space. Using the procedure was taken in [7, 10], we identify the horizon metric

$$ds_h^2 = g_{\varphi\varphi}d\varphi^2 + g_{\theta\theta}d\theta^2|_{r=r_+}, \tag{2.31}$$

with a line element in the cylindrical coordinates

$$ds_3^2 = dz^2 + dR^2 + R^2d\Phi^2, \tag{2.32}$$

where $z = z(\theta)$, $R = R(\theta)$. Then using metric (2.29) and setting $\Phi = \frac{2\pi}{\mu}\varphi$, we have

$$R^2(\theta) = \left(\frac{\mu}{2\pi}\right)g_{\theta\theta}, \tag{2.33}$$

$$\left(\frac{dz(\theta)}{d\theta}\right)^2 = \left(\frac{dR(\theta)}{d\theta}\right)^2 - g_{\varphi\varphi}.$$

The surface of revolution by numerically integrating (2.34) is presented in Fig. 1. That clearly shows a manifold with two punctures at $\theta = 0, \pi$.

Conformal Boundary : Here, we find the conformal boundary of the metric (2.29) by taking conformal factor as l^2/r^2,

$$ds_{bdry}^2 = -dt^2 + 2l\sin^2\theta\,dtd\varphi + \frac{l^2}{\sin^4\theta}d\theta^2. \tag{2.34}$$

By introducing $k = (1-\cos\theta)$, this metric takes the following expression near the pole $\theta = 0$, for small k

$$ds_{bdry}^2 = -dt^2 + 4lk\,dtd\varphi + \frac{l^2}{4k^2}. \tag{2.35}$$



Fig. 1. Horizon embeddings in 4d. Plot shows a 2-dimensional horizons embedded in $\mathbb{R}^3$ as a surface of revolution, displaying topology of a sphere with two punctures. We have set $\mu = 2\pi$, $r_+ = 2$, and $l = 1$ with $s = 0.4$.

That shows an AdS_3 metric written as a Hopf-like fibration over $\mathbb{H}^2$, in which by approaching to the poles, φ becomes a null coordinate. Therefore one can conclude that there is no problem near the poles ($\theta = 0, \pi$), they are not part of the spacetime indeed. It should be noted, one can study the behavior of any outgoing null geodesics that never reach to the $\theta = 0, \pi$ axis in a finite affine parameter [9].

In the extended phase space background, it was shown in [6, 9] that the ultra-spinning general Kerr-AdS black hole solutions exhibit another distinguishable property. In which their entropy for some range of parameter space violate the 'Reverse Isoperimetric Inequality', so they are called as super-entropic black holes. As a further work, we want to explore this approach for our new class of ultra-spinning KK-AdS solutions (2.18).

3 Kerr/CFT description

In this section we explore the Kerr/CFT correspondence for new noncompactness horizon KK-AdS black holes (2.18). The Kerr/CFT correspondence describes a duality between the near horizon states of a 4D extremal kerr black hole and a certain $D = 2$ chiral conformal field theory [17]. This duality has been studied carefully for a large class of black holes, confirming the microscopic entropy of the dual CFT is exactly equivalent to the Bekenstein-Hawking entropy of the bulk geometry.

To start the procedure towards the Kerr/CFT correspondence, at the first we consider the four dimensional ultra-spinning KK-AdS solution (2.29). Then, one need to find the near horizon geometry of the extremal version of the this metric. The extremality conditions can be computed by imposing

vanishing temperature ($T_H|_{r=r_0}=0$), as

$$m=\frac{(l^2+r_0^2)}{2r_0(r_0^2s^2-l^2)},\qquad\qquad s=\frac{l}{r_0}\sqrt{\frac{l^2-3r_0^2}{3l^2-r_0^2}},\tag{3.1}$$

where r_0 denotes the degenerate Killing horizon. Then the function $\hat{\Delta}$ near the horizon is given by

$$\hat{\Delta}=X(r-r_0)^2+\mathcal{O}(r-r_0)^3,$$

where

$$X=\frac{3l^4-2l^2r_0^2+3r_0^4}{2l^2r_0^3}.\tag{3.2}$$

The entropy of the metric (2.29) in the extremal case reads

$$S=\frac{\mu(l^2+r_0^2)}{2r_0}\sqrt{\frac{l^2-r_0^2}{2}}.\tag{3.3}$$

Now, to find the near horizon extremal geometry (NHEG) of the metric (2.29), one can employ the following dimensionless coordinate transformation

$$r=r_0(1+\lambda\hat{r}),\qquad\varphi=\hat{\varphi}+\Omega^0\hat{t},\qquad\hat{t}=\frac{t}{2\pi s_0r_0\lambda},\qquad\hat{\theta}=\theta,\tag{3.4}$$

where the scaling $s_0=\frac{X}{l^2+r_0^2}\sqrt{\frac{2}{l^2-r_0^2}}$, and $\Omega^0=\Omega|_{r=r_0}$. Then, by taking $\lambda\to 0$, we obtain NHEG

$$ds^2 = \sqrt{\hat{H}_4^0}\bigg[\frac{r_0^2+l^2\sin^2\theta}{X}\big(-\hat{r}^2d\hat{t}^2+\frac{d\hat{r}^2}{\hat{r}^2}\big)+\gamma(\theta)\bigg(d\varphi+k\hat{r}d\hat{t}\bigg)^2+F(\theta)d\theta^2\bigg],\tag{3.5}$$

where

$$\begin{aligned}\gamma(\theta) &= \frac{(l^2+r_0^2)^2(l^2-r_0^2)\sin^4\theta}{l^4-r_0^4-2\,l^2r_0^2\sin^2\theta},\qquad F(\theta)=\frac{r_0^2+l^2\cos^2\theta}{\sin^2\theta},\\ \hat{H}_4^0 &= 1+\frac{(l^2-3r_0^2)(l^2+r_0^2)}{2r_0^2(r_0^2+l^2\,\cos^2\theta)}\qquad k=\frac{1}{2\pi T_L}=\frac{2\,l\,r_0}{s_0(l^2+r_0^2)^2}\end{aligned}\tag{3.6}$$

where T_l is the Frolov–Thorne temperature.

The NHEG (3.5) shows a direct product of $\mathrm{AdS}_2\times U(1)$. As expected it contains the well-known sector AdS_2 part which is here written in Poincaré-type coordinates. Appearing of the AdS_2 sector in the NHEG of any regular stationary black holes has been a great appeal in the literature [35–38]. Also it was shown that this class of NHEG by considering a set of consistent boundary conditions admits an enhanced $SL(2,R)\times U(1)$, isometry group [17,39]. Therefore it is argued that there is an algebra of the charges associated to the asymptotic symmetry group that makes a Virasoro algebra with a central extension [39]. This idea supports to existence of a two-dimensional chiral CFT.

One can compute the central charge of the dual quantum field theory using the manner introduced in [20,22]

$$c_{grav}=\frac{3k_i}{2\pi}\int_0^\pi d\theta\sqrt{F(\theta)\gamma(\theta)},\qquad c_{gauge}=0,\tag{3.7}$$

Ultimately, we find the central charge associated to the dual 2D CFT of the NHEG (3.5)

$$c = \mu \frac{6l^3 r_0 (l^2 - r_0^2)(l^2 + r_0^2)^2}{\pi(3l^6 + l^4 r_0^2 + l^2 r_0^4 + 3r_0^6)}. \tag{3.8}$$

We note that, in the extremal case, the vacuum state of the bulk are in a pure state, while the dual 2D CFT has a mixed density matrix by th temperature T_L, is given by Eq. (3.6) .

To ensure that there is a well defined Kerr/CFT duality, we use the Cardy's formula which gives a measure of microstates of any unitary and modular invariance 2D CFT at a large temperature.

$$S = \frac{\pi^2}{3} c_L \, T_L. \tag{3.9}$$

So by assuming that the dual 2D quantum field theory describes a unitary CFT, and using (3.8), (3.6), the microscopic entropy of the dual CFT is computed as

$$S_{CFT} = \frac{\mu(l^2 + r_0^2)}{2r_0} \sqrt{\frac{l^2 - r_0^2}{2}}. \tag{3.10}$$

that is exactly agrees to the Bekenstein-Hawking entropy of the bulk (3.3).

It is worth mentioning that, it was explicitly shown that for some class of charged black holes in [10], the extremality conditions commute with the ultra-spinning limit. Moreover, the commutation of the NHEG and the ultra-spinning limit with each other is examined as well [10,24]. These analysis have done for general multi-spinning Kerr-AdS and two particular classes of gauged supergravity solutions. Now, we explore this statement for the ultra-spinning KK-AdS 4D black hole (2.29).

Extremality under Ultra-spinning limit: We have already derived the extremality conditions for the ultra-spinning version in (3.1). Now, to explore the preserving extremality under ultra-spinning limit, one need to find the extremality conditions of the origin 4D KK-AdS black hole, which is given by

$$m = \frac{(l^2 + r_0^2)(r_0^2 + a^2)}{2r_0(r_0^2 s^2 - l^2)}, \qquad s^2 = \frac{a^2(l^2 - r_0^2 - 3r_0^4 - l^2 r_0^2)}{4m r_0^3}. \tag{3.11}$$

Now, upon taking $a \to l$ limit onto relations above, they precisely agree with the conditions (3.1). Therefore one can confirm again that extremality conditions preserve under ultra-spinning limit.

In order to get ensured about the commutation NHEG and ultra-spinning limit with each other, we should find firstly the near horizon geometry of the 4D KK-AdS black hole in the extremal limit, that is also presented in [42]. Now by applying procedures towards an ultra-spinning version, one can straightforwardly check that the given NHEG exactly gets the metric (3.5).

It should be noted, however the main KK-AdS metric (2.2) and its NHEG that is given in ref. [42] are presented in the asymptotically static frame. To derive the ultra-spinning version of the NHEG, we need not to rewrite NHEG in the ARF. Namely, the near horizon geometry is free of the asymptotic behavior. So, only change coordinate (2.15) and taking limit $a \to l$ onto the NHEG gets the metric (3.5). In the other words, the NHEG of the ultra-spinning version is equivalent to the ultra-spinning version of the NHEG of origin one. Also the sector S^2 of the NHEG (3.5) enjoys a non-compactness structure.

3.1 Kerr/CFT description of general ultraspinning KK-AdS black holes

In this section, we shall investigate the Kerr/CFT correspondence for the general higher-dimensional ($D > 4$) ultra-spinning KK-AdS black holes (2.18).

To find the near horizon geometry of the metric (2.18) in the extremal case, one can find the extremality conditions by imposing $T|_{r=r_0} = 0$ in (2.31), and $\hat{\Delta}|_{r=r_0} = 0$. Then, by taking the following coordinate transformation

$$r = r_0(1+\lambda\hat{r}), \qquad \varphi_j = \hat{\varphi}_j + \Omega_j^0 \hat{t}, \qquad \phi_i = \hat{\phi}_i + \Omega_i^0 \hat{t}, \qquad t = \frac{2\hat{Y}_0}{r_0 \Delta_0'' \lambda}\hat{t}, \qquad \hat{\mu} = \mu. \quad (3.12)$$

and sending $\lambda \to 0$, we derived explicitly the NHEG as

$$\begin{aligned} ds^2 &= \hat{H}_0^{1/(D-2)} \Big[\frac{2\hat{U}_0}{\Delta_0''}\big(-\hat{r}^2 d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2}\big) + \sum_{i,k\neq j}^{N} \tilde{g}_{ik}(d\hat{\phi}_i + k_i\,\hat{r}\,d\hat{t})(d\hat{\phi}_k + k_k\,\hat{r}\,d\hat{t}) \\ &+ \sum_{i}^{N} g_{ij}(d\hat{\phi}_i + k_i\,\hat{r}\,d\hat{t})(d\hat{\varphi}_j + k_j \hat{r}\,d\hat{t}) + d\hat{\Omega}_0^2 \Big], \end{aligned} \quad (3.13)$$

where

$$\begin{aligned} \tilde{g}_{ik\neq j} &= \frac{r_0^2 + a_i^2}{\Xi_i}\mu_i^2 \delta_{ik} + \left(\frac{2m}{\hat{U}_0 \hat{H}^0} + \frac{\hat{\Delta}_0[l^2 s^2 \hat{\Delta}_0 + \hat{U}_0 \hat{H}_0(l^2 - s^2 r_0^2)]}{\hat{H}_0\, c^2\, r_0^{2\epsilon-4} \rho_0^4 \prod_{i\neq j}^{N}(r_0^2 + a_i^2)^2 (c\hat{W} - \sum_{i\neq j}^{N} Z_i \Omega_i^0)} \right) Z_i Z_k, \\ \tilde{g}_{ij} &= (r_0^2 + l^2)\mu_j^2 + \left(\frac{2m}{\hat{U}_0 \hat{H}^0} + \frac{\hat{\Delta}_0[l^2 s^2 \hat{\Delta}_0 + \hat{U}_0 \hat{H}_0(l^2 - s^2 r_0^2)]}{\hat{H}_0\, c^2\, r_0^{2\epsilon-4} \rho_0^4 \prod_{i\neq j}^{N}(r_0^2 + a_i^2)^2 (c\hat{W} - \sum_{i} Z_i \Omega_j^0)} \right) c\, l\, Z_i\, \mu_j^2, \\ Z_i &= \frac{a_i \sqrt{\chi_i}}{\Xi_i}\mu_i^2, \qquad \Delta_0'' = \partial_r \hat{\Delta}|_{r=r_0}, \qquad d\hat{\Omega}_0^2 = d\Omega_s^2|_{r=r_0, \mu=\hat{\mu}}, \end{aligned}$$

and

$$k_{i\neq j} = -\frac{2c\, r_0^{(\epsilon-2)} \rho_0^4 \prod_{i\neq j}^{N}(r_0^2 + a_i^2)}{l\sqrt{l^2 - s^2 r_0^2}} \frac{\partial_{r_0}\Omega_i^0}{\Delta_0''}, \qquad k_j = -\frac{2c\, r_0^{(\epsilon-2)} \rho_0^4 \prod_{i\neq j}^{N}(r_0^2 + a_i^2)}{l\sqrt{l^2 - s^2 r_0^2}} \frac{\partial_{r_0}\Omega_j^0}{\Delta_0''}. \quad (3.14)$$

This result can be adopted in the general form discussed in [22]

$$ds^2 = \Gamma(y)\big(-\hat{r}^2 d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2}\big) + \sum_{\alpha=1}^{n-1} F_\alpha dy_\alpha^2 + \sum_{i,j=1}^{n-1} \tilde{g}_{ij}. \quad (3.15)$$

It has been shown that by imposing consistence boundary conditions, these classes of NHEG admit $[(N-1)/2]$ commuting copies of the Virasoro algebra. In which, each of the Virasoro algebra is generated by the following diffeomorphisms

$$\zeta_{(n)}^{i\neq j} = -e^{in\phi_i}\frac{\partial}{\partial_{\phi_i}} - i\,n\,r\,e^{in\phi_i}\frac{\partial}{\partial_r}, \qquad \zeta_{(n)}^{j} = -e^{in\varphi_j}\frac{\partial}{\partial_{\varphi_j}} - i\,n\,r\,e^{in\varphi_j}\frac{\partial}{\partial_r}, \qquad n = 0, \pm1, \pm2, \ldots \quad (3.16)$$

By considering the same boundary conditions as [20], These diffeomorphisms provide an asymptotic symmetry algebra. That gives N following commuting centerless Virasoro algebra

$$i[\zeta_m^a, \zeta_n^a] = (m-n)\zeta_{m+n}^a. \quad (3.17)$$

The conserved charges $Q_{\zeta_n^i}$ associated to ζ_n^i make a similar algebra with central extension. The Dirac brackets of these conserved charges are attributed to a two dimensional CFT. One can find its associated central charge by [22, 40]

$$c_i = \frac{3\,\mu}{4\pi^2} k_i \int d^{n-1}y_\alpha \Big(det\tilde{g}_{ij} \prod_{\alpha=1}^{n-1} F_\alpha \Big)^{1/2} \int d\phi_1 \dots d\phi_i, \qquad i = 1 \dots n-1, \quad i \neq j. \tag{3.18}$$

Where this relation corresponds to the near-horizon metric (3.13). So, one can rewrite it as follow [40]

$$c_i = \frac{3\mu\, k_i}{4\pi^2} A|_{r_+ = r0}. \tag{3.19}$$

Here A denotes the area of the main black hole. We refer the interested reader to [40] for any details of calculation.

It was introduce two general ansatzs by Mei in [40] that one can explicitly show that, the microscopic entropy of the 2D CFT precisely agrees with the Bekenstein-Hawking entropy of the bulk. The NHEG (3.15) can be cast in the Mei's ansatzs.

Now, by considering the Frolov-Thorne vacuum [17, 41] as a vacuum state for the extremal black hole, one can obtain $[(N-1)/2]$ Frolov–Thorne temperatures associated to each copy of CFTs in a manner given by [20, 22]

$$T_{L\phi_i} = -\frac{\partial T_H/\partial_r}{\partial \Omega_{\phi_i}/\partial_r}|_{r=r_0} = \frac{1}{2\pi\, k_i}, \qquad T_{\varphi_j} = -\frac{\partial T_H/\partial_r}{\partial \Omega_{\varphi_j}/\partial_r}|_{r=r_0} = \frac{1}{2\pi\, k_a}. \tag{3.20}$$

Additionally, one can define the quantity T_e associated to the electric charge as

$$T_e = -\frac{\partial T_H/\partial_r}{\partial \Phi_{\phi_i}/\partial_r}|_{r=r_0}. \tag{3.21}$$

The important relation (3.20) was firstly used in four dimensions by [20]. Then it speculated to be valid for higher dimensions in [22]. It was also proved by [40] that one can generalize the definition (3.20) for all known stationary and axisymmetric extremal black holes. Moreover it was shown that for the ultra-spinning noncompactness horizon solutions, the relation (3.20) may still be valid [24].

The relation (3.20) has an essential role to check the validity of the Kerr/CFT correspondence. Namely, via the Cardy's formula, which gets the entropy of an unitary CFT at temperature T_L, by using the relations (3.19) and (3.20), the microscopic entropy of the conformal field theory can be computed as

$$S_{CFT} = \frac{\pi^2}{3} c_i T_{\phi_i} + \frac{\pi^2}{3} c_j T_{\varphi_j} = \frac{A|_{r=r_0}}{4} = S_{BH}(r_+ = r_0). \tag{3.22}$$

We note, the Mei's ansatzs which gives a well-defined Kerr/CFT correspondence, contain the most stationary and axisymmetric black holes with regular horizon. It was also shown that the general charged KK-AdS black holes (2.2) can be followed by Mei's ansatzs, indicating the Kerr/CFT correspondence is applicable for the general charged KK-AdS black holes as well [**?**].

Now, we emphasis the new ultra-spinning KK-AdS black holes solution (2.18) in all dimensions, despite the noncompactness of their horizons has the form of the Mei's ansatzs. Also as we have discussed in the previous sections the poles appearing in the metric are not part of the spacetime and they do not give rise to a metric singularity. So, for our new obtained regular and noncompactness

horizon black holes (3.15), the result (3.22) shows a precise holographically equivalence between the microscopic entropy of the CFT side and the Bekenstein-Hawking entropy of the bulk. As we have explicitly presented in the previous subsection for four dimensional case.

We also note, it is discussed in [43] that, within the Einstein-Hilbert theories the non-gravitational fields have no contribution to the central charge(s) of the dual CFT. This argument is observed for our new ultra-spinning solutions as well.

Another significance feature of the ultra-spinning KK-AdS black holes is the contribution of the dimensionless parameter μ in the central charges and the entropy of both sides. This parameter which is introduced for compactifying the new azimuthal coordinate φ, should be a limited value to have a finite central charge. Also one can define a new quantity T_μ associated to the chemical potential μ [24].

$$T_\mu = -\frac{\partial T_H/\partial_r}{\partial K_{\phi_i}/\partial_r}|_{r=r_0}, \tag{3.23}$$

Where K is the conjugate variable to the chemical potential , introduced firstly in [6]. Appearing of the chemical potential μ in the central charges and entropy needed to be more studied its role in both CFT and bulk sides.

4 Summery

To generate a new class of charged rotating black hole solutions in Kaluza-Klien theory, we have employed the novel ultra-spinning (super-entropic) limit to the general higher dimensional multi-spinning KK-AdS black holes. Our constructing geometry describes a noncompact horizon solution, that the topology of the event horizon is a sphere with some punctures where arises from the poles of the polar coordinate components. In four dimensional case, we explicitly have shown that these poles are $\theta = 0, \pi$. In the general case the number of punctures were introduced are as many number as the roots of $\mu_j = 0$. We emphasize these poles can be considered as a sort of boundary. Therefore the obtained solution in all dimensions are regular and their entropy is finite.

The new higher dimensional ultra-spinning charged KK-AdS black holes are exact solutions of Einstein-Maxwell dilaton theory in all D dimensions (2.1). We note that, it is impossible to generate a multi ultra-spinning solution by performing more than one direction as an ultra-spinning coordinate.

We have also derived the near horizon geometry of the ultra-spinning KK-AdS solutions in the extremal limit. That shows a direct product of an AdS_2 sector and a S^{d-2} manifold having same punctures as origin black hole. Moreover, for four dimensional case we explicitly have investigated the Kerr/CFT correspondence, by calculating the central charge and the microscopic entropy of the 2D CFT via the Cardy's formula. That our result indicate a precisely agreement between the microscopic and macroscopic entropies.

For the general higher dimensional case, we have utilized the Kerr/CFT correspondence to find the microscopic entropy of their two dimensional chiral CFT side. In the D-dimensional ultra-spinning KK-AdS we are dealing with $[(D-1)/2]$ independent $U(1)$ rotation symmetries that generate $[(D-1)/2]$ copies of Virasoro algebra. We have given a main result for their central charges c_i and c_j, as well as their associated Frolov–Thorne temperatures. Also we discussed how by helping the Cardy's formula, one can check the entropy of the CFT side can be exactly reproduced form the Bekenstein-Hawking entropy.

Acknowledgments

SMN would like to thanks K. Hajian, M. M. Sheikh-Jabbari and M. H. Vahidinaia for useful discussions on the related topics. Also SMN thanks the Institute for Research in Fundamental Sciences (IPM) for hospitality while this project was accomplished.

References

[1] R. Myers and M. Perry, Black Holes in Higher Dimensional Space-Times, Annals Phys. 172 (1986) 304.

[2] R. Emparan and R. C. Myers, Instability of ultra-spinning black holes, JHEP 0309 (2003) 025, [arXiv:hep-th/0308056].

[3] M. M. Caldarelli, R. Emparan, and M. J. Rodriguez, Black Rings in (Anti)-deSitter space, JHEP 0811 (2008) 011, [arXiv:hep-th/0806.1954].

[4] J. Armas and N. Obers, Blackfolds in (Anti)-de Sitter Backgrounds, Phys.Rev. D83 (2011) 084039, [arXiv:hep-th/1012.5081].

[5] M. M. Caldarelli, R. G. Leigh, A. C. Petkou, P. M. Petropoulos, V. Pozzoli, et al., " Vorticity in holographic fluids, PoS CORFU2011 (2011) 076, [arXiv:hep-th/1206.4351].

[6] R. A. Hennigar, D. Kubiznak and R. B. Mann, Super-Entropic Black Holes, Phys. Rev. Lett. 115, no. 3, 031101 (2015), [arXiv:hep-th/1411.4309].

[7] A. Gnecchi, K. Hristov, D. Klemm, C. Toldo, and O. Vaughan, "Rotating black holes in 4d gauged supergravity, JHEP 1401 (2014) 127, [arXiv:hep-th/1311.1795].

[8] D. Klemm, Four-dimensional black holes with unusual horizons, Phys.Rev. D89 (2014) 084007, [arXiv:hep-th/1401.3107].

[9] R. A. Hennigar, D. Kubiznak, R. B. Mann and N. Musoke, Ultraspinning limits and super-entropic black holes, JHEP 1506, 096 (2015), [arXiv:hep-th/1504.07529].

[10] S.M. Noorbakhsh, and M. Ghominejad,"Ultra-spinning gauged supergravity black holes and their Kerr/CFT correspondence", Phys. Rev. D 95, 046002 (2017), [arXiv:1611.02324].

[11] D. Kastor, S. Ray, and J. Traschen, "Enthalpy and the Mechanics of AdS Black Holes", Class.Quant.Grav. 26 (2009) 195011, [arXiv:0904.2765].

[12] M. Cvetic, G. Gibbons, D. Kubiznak, and C. Pope, "Black Hole Enthalpy and an Entropy Inequality for the Thermodynamic Volume", Phys.Rev. D84 (2011) 024037, [arXiv:hep-th/1012.2888].

[13] J. W. York, Phys. Rev. D28 (1983) 2929. W. H. Zurek and K. S. Thorne, Phys. Rev. Lett. 54 (1985) 2171. J. A. Wheeler, A Journey into Gravity and Spacetime Freeman, N.Y. (1990). G. t Hooft, Nucl. Phys. B335 (1990) 138. L. Susskind, L. Thorlacius and R. Uglum, Phys. Rev. D48 (1993) 3743. V. Frolov and I. Novikov, Phys. Rev. D48 (1993) 4545. S. Carlip, Phys. Rev. D51 (1995) 632. M. Cvetic and A. Tseytlin, Phys. Rev. D53 (1996) 5619. F. Larsen and F. Wilczek, Phys. Lett. B375 (1996) 37.

[14] A. Strominger. Black hole entropy from near horizon microstates. JHEP, 02:009, 1998.

[15] S. W. Hawking, M. J. Perry, and A. Stro- minger, Phys. Rev. Lett. 116, 231301 (2016), arXiv:1601.00921 [hep-th] .

[16] H. Afshar, D. Grumiller and M. M. Sheikh-Jabbari, Near Horizon Soft Hairs as Microstates of Three Dimensional Black Holes," [arXiv:hep-th/1607.00009].

[17] M. Guica, T. Hartman, W. Song and A. Strominger, "The Kerr/CFT correspondence," [arXiv:hep-th/0809.4266].

[18] J. D. Brown and M. Henneaux, Comm. Math. Phys. 104 (1986) 207.

[19] J. Cardy, Operator Content Of Two-Dimensional Conformally-Invariant Theories, Nucl. Phys. B270 (1986) 186.

[20] T. Hartman, K. Murata, T. Nishioka and A. Strominger, CFT Duals for Extreme Black Holes, JHEP 0904 (2009) 019 [arXiv:0811.4393 [hep-th]].

[21] H. Lu, J. Mei and C.N. Pope, KerrAdS/CFT correspondence in diverse dimension-arXiv:0811.2225 [hep-th].

[22] D. Chow, M. Cvetic, H. Lu, and C. Pope, Extremal Black Hole/CFT Correspondence in (Gauged) Supergravities, [arXiv:hep-th/0812.2918v2].

[23] G. Compere, K. Murata and T. Nishioka, Central Charges in Extreme Black Hole/CFT Correspondence, JHEP 0905, 077 (2009) [arXiv:0902.1001 [hep-th]]. O. J. C. Dias, H. S. Reall and J. E. Santos, Kerr-CFT and gravitational perturbations, JHEP 0908 (2009) 101, [arXiv:0906.2380 [hep-th]].

[24] M. Sinamuli and R. B. Mann, "Super-Entropic Black Holes and the Kerr-CFT Correspondence", [arXiv:hep-th/1512.07597].

[25] V. Frolov, A. Zelnikov, and U. Bleyer, Ann. Phys. (Leipzig) 499, 371 (1987).

[26] A.N. Aliev, H. Cebeci, and T. Dereli, Phys. Rev. D 77, 124022 (2008).

[27] J. Kunz, D. Maison, F. Navarro-Lerida, and J. Viebahn, Phys. Lett. B639, 95 (2006).

[28] Z.W. Chong, M. Cvetic, H. Lu, and C.N. Pope, Phys. Rev. D 72, 041901 (2005).

[29] D.D.K. Chow, Classical Quantum Gravity 28, 032001 (2011).

[30] S.-Q. Wu, General rotating charged Kaluza-Klein-AdS black holes in higher dimensions, Phys. Rev. D 83 (Jun, 2011) 121502. [arXiv:hep-th/1108.4157v1]

[31] G. W. Gibbons, H. Lu, D. N. Page, and C. N. Pope, Rotating black holes in higher dimensions with a cosmological constant, Phys. Rev. Lett. 93 (2004) 171102, [hep-th/0409155].

[32] G. W. Gibbons, H. Lu, D. N. Page, and C. N. Pope, The general Kerr-de Sitter metrics in all dimensions, J. Geom. Phys. 53 (2005) 49-73, [hep-th/0404008].

[33] J. Kunz, D. Maison, F. Navarro-Lerida and J. Viebahn, Rotating Einstein-Maxwell-Dilaton Black Holes in D Dimensions, Phys. Lett. B 639, 95 (2006) [arXiv:hep-th/0606005].

[34] M. Cvetic, M. J. Duff, P. Hoxha, J. T. Liu, H. Lu, J. X. Lu, R. Martinez-Acosta, C. N. Pope et al., Embedding AdS black holes in ten-dimensions and elevendimensions, Nucl. Phys. B558 (1999) 96-126. [hep-th/9903214].

[35] H. K. Kunduri, J. Lucietti and H. S. Reall, Near-horizon symmetries of extremal black holes, Class. Quant. Grav. 24 (2007) 4169, [arXiv:0705.4214 [hep-th]].

[36] H. K. Kunduri and J. Lucietti, A classification of near-horizon geometries of extremal vacuum black holes, [arXiv:0806.2051 [hep-th]]. H. K. Kunduri and J. Lucietti, Classification of near-horizon geometries of extremal black holes,Living Rev. Rel. 16 (2013) 8, [arXiv:1306.2517 [hep-th]].

[37] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen, and S. P. Trivedi, Rotating attractors, JHEP 0610:058,2006 (JHEP 0610:058,2006) [hep-th/0606244].

[38] M. Johnstone, M.M. Sheikh-Jabbari, J. Simon and H. Yavartanoo, Extremal Black Holes and First Law of Thermodynamics, [arXiv:1305.3157 [hep-th]].

[39] G. Barnich and F. Brandt, Covariant theory of asymptotic symmetries, conservation laws and central charges, Nucl. Phys. B 633, 3 (2002) [arXiv:hep-th/0111246]. G. Barnich and G. Compere, Conserved charges and thermodynamics of the spinning Goedel black hole, Phys. Rev. Lett. 95, 031302 (2005) [arXiv:hep-th/0501102].

[40] Mei, J., The Entropy for General Extremal Black Holes, J. High Energy Phys., 2010(04), (2010). [DOI], [arXiv:1002.1349 [hep-th]].

[41] V.P. Frolov and K.S. Thorne, Renormalized stress-energy tensor near the horizon of a slowly evolving, rotating black hole, Phys. Rev. D39, 2125 (1989).

[42] J. -J. Peng, S.-Q. Wu, Statistical entropies of extremal Kaluza-Klein AdS black holes in arbitrary dimensions, Gen Relativ Gravit (2012) 44:9931005.

[43] T. Hartman, K. Murata, T. Nishioka and A. Strominger, CFT Duals for Extreme Black Holes, JHEP 0904, 019 (2009) [arXiv:0811.4393 [hep-th]].